

Hyperspherical Functions and Linear Representations of the Lorentz Group

V. V. Varlamov

Abstract

A double covering of the proper orthochronous Lorentz group is understood as a complexification of the special unimodular group of second order (a double covering of the 3-dimensional rotation group). In virtue of such an interpretation the matrix elements of finite-dimensional representations of the Lorentz group are studied in terms of hyperspherical functions. Different forms of the hyperspherical functions related to other special functions (hypergeometric series, generalized spherical functions, Jacobi function, Appell functions) are considered. It is shown that there is a close relationship between hyperspherical functions and physical fields defined within $(j, 0) \oplus (0, j)$ representation spaces. Recurrence relations between hyperspherical functions are given.

MSC 2000: 15A66, 22E70, 33C70

PACS 1998: 02.10.Tq, 02.30.Gp

1 Introduction

It is widely accepted that the Lorentz group (a rotation group of the 4-dimensional space-time continuum) is a basic core of relativistic physics. Fields and particles are completely formulated within irreducible representations of the Lorentz or Poincaré group¹ (a motion group of the spacetime continuum). In 1958, Naimark showed that all the finite-dimensional representations of the Lorentz group are equivalent to spinor representations [11]. Due to this fact there is a close relationship between the representations of the Lorentz group and Clifford algebras. At present, it is clear that a spinor structure of physics is deeply rooted in nature. All the physical fields are represented by spinor functions defined on the representation spaces (spinspace²). In essence, the spinor functions present itself a fully algebraic construction within the Clifford algebra theory (see, for example, [18, 19, 20]). However, the Wigner–Mackey scheme of induced representations [26, 10] does not incorporate such an algebraic description. On the other hand, there exists a more useful (for physical applications) method of a generalized regular representation proposed by Vilenkin [22] and based on the theory of special functions. In virtue of a wide use of harmonic analysis many authors considered this method as an alternative to the Wigner–Mackey–Weinberg approach (see, for example, [9]). Moreover, this method naturally incorporates the algebraic description related with the Clifford algebra theory. Namely, it is shown in the present paper (section

¹See Wigner and Weinberg works [26, 25], where physical fields are considered in terms of induced representations of the Poincaré group (see also [10, 4]).

2) that for the spinor functions on the Lorentz group there exists an Euler parametrization which gives rise to hyperspherical functions. In general, the hyperspherical functions are the matrix elements of the finite-dimensional representations of the Lorentz group. It allows to represent these functions by the product of the two hypergeometric functions and later on to relate them with Appell functions.

Further, our consideration based mainly on the so-called Van der Waerden representation of the Lorentz group [24]. Namely, the group $SL(2, \mathbb{C})$ (a double covering of the proper orthochronous Lorentz group $\mathfrak{G}_+$) is understood as a complexification of the group $SU(2)$, $SL(2, \mathbb{C}) \sim \text{complex}(SU(2))$. In a sense, it allows to represent the group $SL(2, \mathbb{C})$ by a product $SU(2) \otimes SU(2)^3$ as it done by Ryder in his textbook [14]. In connection with this, the hyperspherical functions should be considered as a complexification of generalized spherical functions for the group $SU(2)$ introduced by Gel'fand and Shapiro in 1952 [7].

The paper is organized as follows. In the section 2 we briefly discuss the basic notions concerning representations of the Lorentz group, the group $SU(2)$, and also we consider their descriptions within Clifford algebra framework. In the section 3 we introduce the main objects of our study, namely, hyperspherical functions. Different forms of hyperspherical functions are given. Recurrence relations between hyperspherical functions, presented one of the main results of the paper, are studied in the section 5.

2 Preliminaries

In this section we will consider some basic facts concerning the Lorentz group and its representations.

2.1 Van der Waerden representation of the Lorentz group

Let $\mathfrak{g} \rightarrow T_{\mathfrak{g}}$ be an arbitrary linear representation of the proper orthochronous Lorentz group $\mathfrak{G}_+$ and let $A_i(t) = T_{a_i(t)}$ be an infinitesimal operator corresponded the rotation $a_i(t) \in \mathfrak{G}_+$. Analogously, we have $B_i(t) = T_{b_i(t)}$, where $b_i(t) \in \mathfrak{G}_+$ is a hyperbolic rotation. The operators

³Moreover, in the works [2, 5] the Lorentz group is represented by a product $SU_R(2) \otimes SU_L(2)$, and spinors

$$\psi(p^\mu) = \begin{pmatrix} \phi_R(p^\mu) \\ \phi_L(p^\mu) \end{pmatrix}$$

($\phi_R(p^\mu)$ and $\phi_L(p^\mu)$ are the right- and left-handed spinors) are transformed within $(j, 0) \oplus (0, j)$ representation space.

A_i and B_i satisfy the following commutation relations⁴:

$$\begin{aligned}
[A_1, A_2] &= A_3, & [A_2, A_3] &= A_1, & [A_3, A_1] &= A_2, \\
[B_1, B_2] &= -A_3, & [B_2, B_3] &= -A_1, & [B_3, B_1] &= -A_2, \\
[A_1, B_1] &= 0, & [A_2, B_2] &= 0, & [A_3, B_3] &= 0, \\
[A_1, B_2] &= B_3, & [A_1, B_3] &= -B_2, \\
[A_2, B_3] &= B_1, & [A_2, B_1] &= -B_3, \\
[A_3, B_1] &= B_2, & [A_3, B_2] &= -B_1.
\end{aligned} \tag{1}$$

Let us consider the operators

$$\begin{aligned}
X_k &= \frac{1}{2}(A_k + iB_k), & Y_k &= \frac{1}{2}(A_k - iB_k), \\
(k &= 1, 2, 3).
\end{aligned} \tag{2}$$

Using the relations (1) we find that

$$\begin{aligned}
[X_1, X_2] &= X_3, & [X_2, X_3] &= X_1, & [X_3, X_1] &= X_2, \\
[Y_1, Y_2] &= Y_3, & [Y_2, Y_3] &= Y_1, & [Y_3, Y_1] &= Y_2, \\
[X_k, Y_l] &= 0, & (k, l &= 1, 2, 3).
\end{aligned} \tag{3}$$

Further, taking

$$\begin{aligned}
X_+ &= X_1 + iX_2, & X_- &= X_1 - iX_2, \\
Y_+ &= Y_1 + iY_2, & Y_- &= Y_1 - iY_2
\end{aligned} \tag{4}$$

we see that in virtue of commutativity of the relations (3) a space of an irreducible finite-dimensional representation of the group $\mathfrak{G}_+$ can be stretched on the totality of $(2l+1)(2l'+1)$ basis vectors $|l, m; l', m'\rangle$, where l, m, l', m' are integer or half-integer numbers, $-l \leq m \leq l$, $-l' \leq m' \leq l'$. Therefore,

$$\begin{aligned}
X_- |l, m; l', m'\rangle &= \sqrt{(l' + m')(l' - m' + 1)} |l, m; l', m' - 1\rangle \quad (m' > -l'), \\
X_+ |l, m; l', m'\rangle &= \sqrt{(l' - m')(l' + m' + 1)} |l, m; l', m' + 1\rangle \quad (m' < l'), \\
X_3 |l, m; l', m'\rangle &= m' |l, m; l', m'\rangle, \\
Y_- |l, m; l', m'\rangle &= \sqrt{(l + m)(l - m + 1)} |l, m - 1; l', m'\rangle \quad (m > -l) \\
Y_+ |l, m; l', m'\rangle &= \sqrt{(l - m)(l + m + 1)} |l, m + 1; l', m'\rangle \quad (m < l), \\
Y_3 |l, m; l', m'\rangle &= m |l, m; l', m'\rangle.
\end{aligned} \tag{5}$$

From the relations (3) it follows that each of the sets of infinitesimal operators X and Y generates the group $SU(2)$ and these two groups commute with each other. Thus, from the relations (3) and (5) it follows that the group $\mathfrak{G}_+$, in essence, is equivalent to the group $SU(2) \otimes SU(2)$. In contrast to the Gel'fand–Naimark representation for the Lorentz group [8, 11] which does not find a broad application in physics, a representation (5) is a most useful in theoretical physics (see, for example, [1, 15, 13, 14]). This representation for the Lorentz group was firstly given by Van der Waerden in his brilliant book [24].

⁴Denoting $I^{23} = A_1$, $I^{31} = A_2$, $I^{12} = A_3$, and $I^{01} = B_1$, $I^{02} = B_2$, $I^{03} = B_3$ we can write the relations (1) in a more compact form:

$$[I^{\mu\nu}, I^{\lambda\rho}] = \delta_{\mu\rho} I^{\lambda\nu} + \delta_{\nu\lambda} I^{\mu\rho} - \delta_{\nu\rho} I^{\mu\lambda} - \delta_{\mu\lambda} I^{\nu\rho}.$$

2.2 Clifford algebras and linear representations of the Lorentz group. Spinor functions

As known, a double covering of the proper orthochronous Lorentz group $\mathfrak{G}_+$, the group $SL(2, \mathbb{C})$, is isomorphic to the Clifford–Lipschitz group $\mathbf{Spin}_+(1, 3)$, which, in its turn, is fully defined within a biquaternion algebra $\mathbb{C}_2$, since

$$\mathbf{Spin}_+(1, 3) \simeq \left\{ \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \in \mathbb{C}_2 : \det \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} = 1 \right\} = SL(2, \mathbb{C}).$$

Thus, a fundamental representation of the group $\mathfrak{G}_+$ is realized in a spinspace $\mathbb{S}_2$. The spinspace $\mathbb{S}_2$ is a complexification of the minimal left ideal of the algebra $\mathbb{C}_2$: $\mathbb{S}_2 = \mathbb{C} \otimes I_{2,0} = \mathbb{C} \otimes \mathcal{O}_{2,0}e_{20}$ or $\mathbb{S}_2 = \mathbb{C} \otimes I_{1,1} = \mathbb{C} \otimes \mathcal{O}_{1,1}e_{11}$ ($\mathbb{C} \otimes I_{0,2} = \mathbb{C} \otimes \mathcal{O}_{0,2}e_{02}$), where $\mathcal{O}_{p,q}$ ($p+q=2$) is a real subalgebra of $\mathbb{C}_2$, $I_{p,q}$ is the minimal left ideal of the algebra $\mathcal{O}_{p,q}$, e_{pq} is a primitive idempotent.

Further, let $\mathbb{C}_2^*$ be the biquaternion algebra, in which all the coefficients are complex conjugate to the coefficients of the algebra $\mathbb{C}_2$. The algebra $\mathbb{C}_2^*$ is obtained from $\mathbb{C}_2$ under action of the automorphism $\mathcal{A} \rightarrow \mathcal{A}^*$ (involution) or the antiautomorphism $\mathcal{A} \rightarrow \tilde{\mathcal{A}}$ (reversal), where $\mathcal{A} \in \mathbb{C}_2$ (see [16, 17]). Let us compose a tensor product of k algebras $\mathbb{C}_2$ and r algebras $\mathbb{C}_2^*$:

$$\mathbb{C}_2 \otimes \mathbb{C}_2 \otimes \cdots \otimes \mathbb{C}_2 \otimes \mathbb{C}_2^* \otimes \mathbb{C}_2^* \otimes \cdots \otimes \mathbb{C}_2^* \simeq \mathbb{C}_{2k} \otimes \mathbb{C}_{2r}^*. \quad (6)$$

The tensor product (6) induces a spinspace

$$\mathbb{S}_2 \otimes \mathbb{S}_2 \otimes \cdots \otimes \mathbb{S}_2 \otimes \mathbb{S}_2^* \otimes \mathbb{S}_2^* \otimes \cdots \otimes \mathbb{S}_2^* = \mathbb{S}_{2k+r} \quad (7)$$

with ‘vectors’ (spintensors) of the form

$$\xi^{\alpha_1 \alpha_2 \cdots \alpha_k \dot{\alpha}_1 \dot{\alpha}_2 \cdots \dot{\alpha}_r} = \sum \xi^{\alpha_1} \otimes \xi^{\alpha_2} \otimes \cdots \otimes \xi^{\alpha_k} \otimes \xi^{\dot{\alpha}_1} \otimes \xi^{\dot{\alpha}_2} \otimes \cdots \otimes \xi^{\dot{\alpha}_r}. \quad (8)$$

The full representation space $\mathbb{S}_{2k+r}$ contains both symmetric and antisymmetric spintensors (8). Usually, at the definition of irreducible finite-dimensional representations of the Lorentz group physicists confined to a subspace of symmetric spintensors $\text{Sym}(k, r) \subset \mathbb{S}_{2k+r}$. Dimension of $\text{Sym}(k, r)$ is equal to $(k+1)(r+1)$ or $(2l+1)(2l'+1)$ at $l = \frac{k}{2}$, $l' = \frac{r}{2}$. It is easy to see that the space $\text{Sym}(k, r)$ is a space of van der Waerden representation (5). The space $\text{Sym}(k, r)$ can be considered as a space of polynomials

$$p(z_0, z_1, \bar{z}_0, \bar{z}_1) = \sum_{\substack{(\alpha_1, \dots, \alpha_k) \\ (\dot{\alpha}_1, \dots, \dot{\alpha}_r)}} \frac{1}{k! r!} a^{\alpha_1 \cdots \alpha_k \dot{\alpha}_1 \cdots \dot{\alpha}_r} z_{\alpha_1} \cdots z_{\alpha_k} \bar{z}_{\dot{\alpha}_1} \cdots \bar{z}_{\dot{\alpha}_r} \quad (\alpha_i, \dot{\alpha}_i = 0, 1), \quad (9)$$

where the numbers $a^{\alpha_1 \cdots \alpha_k \dot{\alpha}_1 \cdots \dot{\alpha}_r}$ are unaffected at the permutations of indices. Some applications of the functions (9) contained in [21, 9].

The representation of the group $\mathfrak{G}_+$ in the space $\text{Sym}(k, r)$ has a form

$$\begin{aligned} T_g q(\zeta, \bar{\zeta}) &= \frac{1}{z_1^k \bar{z}_1^r} T_g \left[z_1^k \bar{z}_1^r q \left(\frac{z_0}{z_1}, \frac{\bar{z}_0}{\bar{z}_1} \right) \right] = \\ &= (\gamma \zeta + \delta)^k (\gamma^* \bar{\zeta} + \delta^*)^r q \left(\frac{\alpha \zeta + \beta}{\gamma \zeta + \delta}, \frac{\alpha^* \bar{\zeta} + \beta^*}{\gamma^* \bar{\zeta} + \delta^*} \right) \end{aligned} \quad (10)$$

where

$$\zeta = \frac{z_0}{z_1}, \quad \bar{\zeta} = \frac{\bar{z}_0}{\bar{z}_1}.$$

It is easy to see that for the group $SU(2) \subset SL(2, \mathbb{C})$ the formulae (9) and (10) reduce to the following

$$p(z_0, z_1) = \sum_{(\alpha_1, \dots, \alpha_k)} \frac{1}{k!} a^{\alpha_1 \dots \alpha_k} z_{\alpha_1} \dots z_{\alpha_k}, \quad (11)$$

$$T_g q(\zeta) = (\gamma\zeta + \delta)^k q\left(\frac{\alpha\zeta + \beta}{\gamma\zeta + \delta}\right), \quad (12)$$

and the representation space $\text{Sym}(k, r)$ reduces to $\text{Sym}(k, 0)$. One-parameter subgroups of $SU(2)$ are defined by the matrices

$$a_1(t) = \begin{pmatrix} \cos \frac{t}{2} & i \sin \frac{t}{2} \\ i \sin \frac{t}{2} & \cos \frac{t}{2} \end{pmatrix}, \quad a_2(t) = \begin{pmatrix} \cos \frac{t}{2} & -\sin \frac{t}{2} \\ \sin \frac{t}{2} & \cos \frac{t}{2} \end{pmatrix}, \quad a_3(t) = \begin{pmatrix} e^{\frac{it}{2}} & 0 \\ 0 & e^{-\frac{it}{2}} \end{pmatrix}. \quad (13)$$

An arbitrary matrix $u \in SU(2)$ written via Euler angles has a form

$$u = \begin{pmatrix} \alpha & \beta \\ -\beta^* & \alpha^* \end{pmatrix} = \begin{pmatrix} \cos \frac{\theta}{2} e^{\frac{i(\varphi+\psi)}{2}} & i \sin \frac{\theta}{2} e^{\frac{i(\varphi-\psi)}{2}} \\ i \sin \frac{\theta}{2} e^{\frac{i(\psi-\varphi)}{2}} & \cos \frac{\theta}{2} e^{-\frac{i(\varphi+\psi)}{2}} \end{pmatrix}, \quad (14)$$

where $0 \leq \varphi < 2\pi$, $0 < \theta < \pi$, $-2\pi \leq \psi < 2\pi$, $\det u = 1$. Hence it follows that $|\alpha| = \cos \frac{\theta}{2}$, $|\beta| = \sin \frac{\theta}{2}$ and

$$\cos \theta = 2|\alpha|^2 - 1, \quad (15)$$

$$e^{i\varphi} = -\frac{\alpha\beta i}{|\alpha||\beta|}, \quad (16)$$

$$e^{\frac{i\psi}{2}} = \frac{\alpha e^{-\frac{i\varphi}{2}}}{|\alpha|}. \quad (17)$$

Diagonal matrices $\begin{pmatrix} e^{\frac{i\varphi}{2}} & 0 \\ 0 & e^{-\frac{i\varphi}{2}} \end{pmatrix}$ form one-parameter subgroup in the group $SU(2)$. Therefore, each matrix $u \in SU(2)$ belongs to a bilateral adjacency class contained the matrix

$$\begin{pmatrix} \cos \frac{\theta}{2} & i \sin \frac{\theta}{2} \\ i \sin \frac{\theta}{2} & \cos \frac{\theta}{2} \end{pmatrix}$$

The matrix element $t_{mn}^l = e^{-i(m\varphi+n\psi)} \langle T_l(\theta) \psi_n, \psi_m \rangle$ of the group $SU(2)$ in the polynomial basis

$$\zeta^{l-n}$$

where

$$T_l(\theta)\psi(\zeta) = \left(i \sin \frac{\theta}{2} \zeta + \cos \frac{\theta}{2}\right)^{2l} \psi \left(\frac{\cos \frac{\theta}{2} \zeta + i \sin \frac{\theta}{2}}{i \sin \frac{\theta}{2} \zeta + \cos \frac{\theta}{2}} \right),$$

has a form

$$\begin{aligned} t_{mn}^l(g) &= e^{-i(m\varphi+n\psi)} \langle T_l(\theta)\psi_n, \psi_m \rangle = \\ &= \frac{e^{-i(m\varphi+n\psi)} \langle T_l(\theta)\zeta^{l-n}\zeta^{l-m} \rangle}{\sqrt{\Gamma(l-m+1)\Gamma(l+m+1)\Gamma(l-n+1)\Gamma(l+n+1)}} = \\ &= e^{-i(m\varphi+n\psi)} i^{m-n} \sqrt{\Gamma(l-m+1)\Gamma(l+m+1)\Gamma(l-n+1)\Gamma(l+n+1)} \times \\ &\quad \cos^{2l} \frac{\theta}{2} \tan^{m-n} \frac{\theta}{2} \sum_{j=\max(0, n-m)}^{\min(l-n, l+n)} \frac{i^{2j} \tan^{2j} \frac{\theta}{2}}{\Gamma(j+1)\Gamma(l-m-j+1)\Gamma(l+n-j+1)\Gamma(m-n+j+1)}. \end{aligned} \quad (18)$$

Further, using the formula

$${}_2F_1(\alpha, \beta; \gamma; z) = \frac{\Gamma(\gamma)}{\Gamma(\alpha)\Gamma(\beta)} \sum_{k \geq 0} \frac{\Gamma(\alpha+k)\Gamma(\beta+k)}{\Gamma(\gamma+k)} \frac{z^k}{k!} \quad (19)$$

we express the matrix element (18) via the hypergeometric function:

$$\begin{aligned} t_{mn}^l(g) &= \frac{i^{m-n} e^{-i(m\varphi+n\psi)}}{\Gamma(m-n+1)} \sqrt{\frac{\Gamma(l+m+1)\Gamma(l-n+1)}{\Gamma(l-m+1)\Gamma(l+n+1)}} \times \\ &\quad \cos^{2l} \frac{\theta}{2} \tan^{m-n} \frac{\theta}{2} {}_2F_1(m-l+1, 1-l-n, m-n+1; i^2 \tan^2 \frac{\theta}{2}), \end{aligned} \quad (20)$$

where $m \geq n$. At $m < n$ in the right part of (20) it needs to replace m and n by $-m$ and $-n$, respectively. Since l, m and n are finite numbers, then the hypergeometric series is interrupted.

Further, replacing in the one-parameter subgroups (13) the parameter t by $-it$ we obtain

$$b_1(t) = \begin{pmatrix} \cosh \frac{t}{2} & \sinh \frac{t}{2} \\ \sinh \frac{t}{2} & \cosh \frac{t}{2} \end{pmatrix}, \quad b_2(t) = \begin{pmatrix} \cosh \frac{t}{2} & i \sinh \frac{t}{2} \\ -i \sinh \frac{t}{2} & \cosh \frac{t}{2} \end{pmatrix}, \quad b_3(t) = \begin{pmatrix} e^{\frac{t}{2}} & 0 \\ 0 & e^{-\frac{t}{2}} \end{pmatrix}. \quad (21)$$

These subgroups correspond to hyperbolic rotations.

3 Hyperspherical functions

The group $SL(2, \mathbb{C})$ of all complex matrices

$$\begin{pmatrix} \alpha & \beta \end{pmatrix}$$

of 2-nd order with the determinant $\alpha\delta - \gamma\beta = 1$, is a *complexification* of the group $SU(2)$. The group $SU(2)$ is one of the real forms of $SL(2, \mathbb{C})$. The transition from $SU(2)$ to $SL(2, \mathbb{C})$ is realized via the complexification of three real parameters φ, θ, ψ (Euler angles). Let $\theta^c = \theta - i\tau, \varphi^c = \varphi - i\epsilon, \psi^c = \psi - i\varepsilon$ be complex Euler angles, where

$$\begin{aligned} 0 &\leq \operatorname{Re} \theta^c = \theta \leq \pi, & 0 &< \operatorname{Im} \theta^c = \tau < \infty, \\ 0 &\leq \operatorname{Re} \varphi^c = \varphi < 2\pi, & 0 &\leq \operatorname{Im} \varphi^c = \epsilon \leq 2\pi, \\ -2\pi &\leq \operatorname{Re} \psi^c = \psi < 2\pi, & -2\pi &\leq \operatorname{Im} \psi^c = \varepsilon < 2\pi. \end{aligned}$$

Replacing in (14) the angles φ, θ, ψ by the complex angles $\varphi^c, \theta^c, \psi^c$ we come to the following matrix

$$\begin{aligned} \mathfrak{g} &= \begin{pmatrix} \cos \frac{\theta^c}{2} e^{\frac{i(\varphi^c + \psi^c)}{2}} & i \sin \frac{\theta^c}{2} e^{\frac{i(\varphi^c - \psi^c)}{2}} \\ i \sin \frac{\theta^c}{2} e^{\frac{i(\psi^c - \varphi^c)}{2}} & \cos \frac{\theta^c}{2} e^{-\frac{i(\varphi^c + \psi^c)}{2}} \end{pmatrix} = \\ &= \begin{pmatrix} \left(\cos \frac{\theta}{2} \cosh \frac{\tau}{2} + i \sin \frac{\theta}{2} \sinh \frac{\tau}{2} \right) e^{\frac{\epsilon + \varepsilon + i(\varphi + \psi)}{2}} & \left(\cos \frac{\theta}{2} \sinh \frac{\tau}{2} + i \sin \frac{\theta}{2} \cosh \frac{\tau}{2} \right) e^{\frac{\epsilon - \varepsilon + i(\varphi - \psi)}{2}} \\ \left(\cos \frac{\theta}{2} \sinh \frac{\tau}{2} + i \sin \frac{\theta}{2} \cosh \frac{\tau}{2} \right) e^{\frac{\varepsilon - \epsilon + i(\psi - \varphi)}{2}} & \left(\cos \frac{\theta}{2} \cosh \frac{\tau}{2} + i \sin \frac{\theta}{2} \sinh \frac{\tau}{2} \right) e^{\frac{-\epsilon - \varepsilon - i(\varphi + \psi)}{2}} \end{pmatrix}, \end{aligned} \quad (22)$$

since $\cos \frac{1}{2}(\theta - i\tau) = \cos \frac{\theta}{2} \cosh \frac{\tau}{2} + i \sin \frac{\theta}{2} \sinh \frac{\tau}{2}$, and $\sin \frac{1}{2}(\theta - i\tau) = \sin \frac{\theta}{2} \cosh \frac{\tau}{2} - i \cos \frac{\theta}{2} \sinh \frac{\tau}{2}$. It is easy to verify that the matrix (22) coincides with a matrix of the fundamental representation of the group $SL(2, \mathbb{C})$ (in Euler parametrization):

$$\begin{aligned} \mathfrak{g}(\varphi, \epsilon, \theta, \tau, \psi, \varepsilon) &= \\ &= \begin{pmatrix} e^{i\frac{\varphi}{2}} & 0 \\ 0 & e^{-i\frac{\varphi}{2}} \end{pmatrix} \begin{pmatrix} e^{\frac{\epsilon}{2}} & 0 \\ 0 & e^{-\frac{\epsilon}{2}} \end{pmatrix} \begin{pmatrix} \cos \frac{\theta}{2} & i \sin \frac{\theta}{2} \\ i \sin \frac{\theta}{2} & \cos \frac{\theta}{2} \end{pmatrix} \begin{pmatrix} \cosh \frac{\tau}{2} & \sinh \frac{\tau}{2} \\ \sinh \frac{\tau}{2} & \cosh \frac{\tau}{2} \end{pmatrix} \begin{pmatrix} e^{i\frac{\psi}{2}} & 0 \\ 0 & e^{-i\frac{\psi}{2}} \end{pmatrix} \begin{pmatrix} e^{\frac{\varepsilon}{2}} & 0 \\ 0 & e^{-\frac{\varepsilon}{2}} \end{pmatrix}. \end{aligned} \quad (23)$$

The matrix element $t_{mn}^l = e^{-m(\epsilon + i\varphi) - n(\varepsilon + i\psi)} \langle T_l(\theta, \tau) \psi_\lambda, \psi_{\dot{\lambda}} \rangle$ of the finite-dimensional representation of $SL(2, \mathbb{C})$ at $l = l'$ in the polynomial basis

$$\psi_\lambda(\zeta, \bar{\zeta}) = \frac{\zeta^{l-n} \bar{\zeta}^{l-m}}{\sqrt{\Gamma(l-n+1)\Gamma(l+n+1)\Gamma(l-m+1)\Gamma(l+m+1)}},$$

has a form

$$\begin{aligned}
t_{mn}^l(\mathfrak{g}) &= e^{-m(\epsilon+i\varphi)-n(\epsilon+i\psi)} Z_{mn}^l = e^{-m(\epsilon+i\varphi)-n(\epsilon+i\psi)} \times \\
&\sum_{k=-l}^l i^{m-k} \sqrt{\Gamma(l-m+1)\Gamma(l+m+1)\Gamma(l-k+1)\Gamma(l+k+1)} \cos^{2l} \frac{\theta}{2} \tan^{m-k} \frac{\theta}{2} \times \\
&\sum_{j=\max(0,k-m)}^{\min(l-m,l+k)} \frac{i^{2j} \tan^{2j} \frac{\theta}{2}}{\Gamma(j+1)\Gamma(l-m-j+1)\Gamma(l+k-j+1)\Gamma(m-k+j+1)} \times \\
&\sqrt{\Gamma(l-n+1)\Gamma(l+n+1)\Gamma(l-k+1)\Gamma(l+k+1)} \cosh^{2l} \frac{\tau}{2} \tanh^{n-k} \frac{\tau}{2} \times \\
&\sum_{s=\max(0,k-n)}^{\min(l-n,l+k)} \frac{\tanh^{2s} \frac{\tau}{2}}{\Gamma(s+1)\Gamma(l-n-s+1)\Gamma(l+k-s+1)\Gamma(n-k+s+1)}. \quad (24)
\end{aligned}$$

We will call the functions Z_{mn}^l in (24) as *hyperspherical functions*. Using (19) we can write the hyperspherical functions Z_{mn}^l via the hypergeometric series:

$$\begin{aligned}
Z_{mn}^l &= \cos^{2l} \frac{\theta}{2} \cosh^{2l} \frac{\tau}{2} \sum_{k=-l}^l i^{m-k} \tan^{m-k} \frac{\theta}{2} \tanh^{n-k} \frac{\tau}{2} \times \\
&{}_2F_1(m-l+1, 1-l-k; m-k+1; i^2 \tan^2 \frac{\theta}{2}) {}_2F_1(n-l+1, 1-l-k; n-k+1; \tanh^2 \frac{\tau}{2}) \quad (25)
\end{aligned}$$

Therefore, matrix elements can be expressed by means of the function (*a generalized hyperspherical function*)

$$\mathfrak{M}_{mn}^l(\mathfrak{g}) = e^{-m(\epsilon+i\varphi)} Z_{mn}^l e^{-n(\epsilon+i\psi)}, \quad (26)$$

where

$$Z_{mn}^l = \sum_{k=-l}^l P_{mk}^l(\cos \theta) \mathfrak{P}_{kn}^l(\cosh \tau), \quad (27)$$

here $P_{mn}^l(\cos \theta)$ is a generalized spherical function on the group $SU(2)$ (see [8]), and $\mathfrak{P}_{mn}^l$ is an analog of the generalized spherical function for the group $QU(2)$ (so-called Jacobi function [22]). $QU(2)$ is a group of quasiunitary unimodular matrices of second order. As well as the group $SU(2)$ the group $QU(2)$ is one of the real forms of $SL(2, \mathbb{C})$ ($QU(2)$ is noncompact).

Further, from (25) we see that the function Z_{mn}^l depends on two variables θ and τ . Therefore, in particular cases we can express the hyperspherical functions Z_{mn}^l via Appell functions F_1 – F_4 (hypergeometric series of two variables [3, 6]). However, a consideration of the relations with Appell functions comes beyond the framework of this paper and we will consider it in a separate work.

3.1 Matrices $T_l(\mathfrak{g})$

Using the formula (24) let us find explicit expressions for the matrices $T_l(\mathfrak{g})$ of the finite-dimensional representations of $\mathfrak{G}_+$ at $l = 0, \frac{1}{2}, 1$:

$$T_0(\theta, \tau) = 1,$$

$$T_{\frac{1}{2}}(\theta, \tau) = \begin{pmatrix} Z_{-\frac{1}{2}-\frac{1}{2}}^{\frac{1}{2}} & Z_{\frac{1}{2}-\frac{1}{2}}^{\frac{1}{2}} \\ Z_{-\frac{1}{2}\frac{1}{2}}^{\frac{1}{2}} & Z_{\frac{1}{2}\frac{1}{2}}^{\frac{1}{2}} \end{pmatrix} = \begin{pmatrix} \cos \frac{\theta}{2} \cosh \frac{\tau}{2} + i \sin \frac{\theta}{2} \sinh \frac{\tau}{2} & \cos \frac{\theta}{2} \sinh \frac{\tau}{2} + i \sin \frac{\theta}{2} \cosh \frac{\tau}{2} \\ \cos \frac{\theta}{2} \sinh \frac{\tau}{2} + i \sin \frac{\theta}{2} \cosh \frac{\tau}{2} & \cos \frac{\theta}{2} \cosh \frac{\tau}{2} + i \sin \frac{\theta}{2} \sinh \frac{\tau}{2} \end{pmatrix}, \quad (28)$$

$$T_1(\theta, \tau) = \begin{pmatrix} Z_{-1-1}^1 & Z_{-10}^1 & Z_{-11}^1 \\ Z_{0-1}^1 & Z_{00}^1 & Z_{01}^1 \\ Z_{1-1}^1 & Z_{10}^1 & Z_{11}^1 \end{pmatrix} = \begin{pmatrix} \cos^2 \frac{\theta}{2} \cosh^2 \frac{\tau}{2} + \frac{i \sin \theta \sinh \tau}{2} - \sin^2 \frac{\theta}{2} \sinh^2 \frac{\tau}{2} & \frac{1}{\sqrt{2}}(\cos \theta \sinh \tau + i \sin \theta \cosh \tau) & \cos^2 \frac{\theta}{2} \sinh^2 \frac{\tau}{2} + \frac{i \sin \theta \sinh \tau}{2} - \sin^2 \frac{\theta}{2} \cosh^2 \frac{\tau}{2} \\ \frac{1}{\sqrt{2}}(\cos \theta \sinh \tau + i \sin \theta \cosh \tau) & \cos \theta \cosh \tau + i \sin \theta \sinh \tau & \frac{1}{\sqrt{2}}(\cos \theta \sinh \tau + i \sin \theta \cosh \tau) \\ \cos^2 \frac{\theta}{2} \sinh^2 \frac{\tau}{2} + \frac{i \sin \theta \sinh \tau}{2} - \sin^2 \frac{\theta}{2} \cosh^2 \frac{\tau}{2} & \frac{1}{\sqrt{2}}(\cos \theta \sinh \tau + i \sin \theta \cosh \tau) & \cos^2 \frac{\theta}{2} \cosh^2 \frac{\tau}{2} + \frac{i \sin \theta \sinh \tau}{2} - \sin^2 \frac{\theta}{2} \sinh^2 \frac{\tau}{2} \end{pmatrix}. \quad (29)$$

3.2 Addition Theorem

Let $\mathfrak{g} = \mathfrak{g}_1 \mathfrak{g}_2$ be the product of two matrices $\mathfrak{g}_1, \mathfrak{g}_2 \in SL(2, \mathbb{C})$. Let us denote the Euler angles of the matrix $\mathfrak{g}$ via $\varphi^c, \theta^c, \psi^c$, the matrix $\mathfrak{g}_1$ via $\varphi_1^c, \theta_1^c, \psi_1^c$ and the matrix $\mathfrak{g}_2$ via $\varphi_2^c, \theta_2^c, \psi_2^c$. Expressing now the Euler angles of the matrix $\mathfrak{g}$ via the Euler angles of the factors $\mathfrak{g}_1, \mathfrak{g}_2$ we consider at first the particular case $\varphi_1^c = \psi_1^c = \psi_2^c = 0$:

$$\mathfrak{g} = \begin{pmatrix} \cos \frac{\theta_1^c}{2} & i \sin \frac{\theta_1^c}{2} \\ i \sin \frac{\theta_1^c}{2} & \cos \frac{\theta_1^c}{2} \end{pmatrix} \begin{pmatrix} \cos \frac{\theta_2^c}{2} e^{\frac{i\varphi_2^c}{2}} & i \sin \frac{\theta_2^c}{2} e^{\frac{i\varphi_2^c}{2}} \\ i \sin \frac{\theta_2^c}{2} e^{-\frac{i\varphi_2^c}{2}} & \cos \frac{\theta_2^c}{2} e^{-\frac{i\varphi_2^c}{2}} \end{pmatrix}.$$

Multiplying the matrices in the right part of this equality and using a complex analog of the formulae (15)–(17) we obtain

$$\cos \theta^c = \cos \theta_1^c \cos \theta_2^c - \sin \theta_1^c \sin \theta_2^c \cos \varphi_2^c, \quad (30)$$

$$e^{i\varphi^c} = \frac{\sin \theta_1^c \cos \theta_2^c + \cos \theta_1^c \sin \theta_2^c \cos \varphi_2^c + i \sin \theta_2^c \sin \varphi_2^c}{\sin \theta^c}, \quad (31)$$

$$e^{\frac{i(\varphi^c + \psi^c)}{2}} = \frac{\cos \frac{\theta_1^c}{2} \cos \frac{\theta_2^c}{2} e^{\frac{i\varphi_2^c}{2}} - \sin \frac{\theta_1^c}{2} \sin \frac{\theta_2^c}{2} e^{-\frac{i\varphi_2^c}{2}}}{\sin \frac{\theta^c}{2}}. \quad (32)$$

It is not difficult to obtain a general case. Indeed, in virtue of (23) the matrix $\mathbf{g} \in SL(2, \mathbb{C})$ admits a representation

$$\begin{aligned} \mathbf{g}(\varphi^c, \theta^c, \psi^c) &= \begin{pmatrix} e^{\frac{i\varphi^c}{2}} & 0 \\ 0 & e^{-\frac{i\varphi^c}{2}} \end{pmatrix} \begin{pmatrix} \cos \frac{\theta^c}{2} & i \sin \frac{\theta^c}{2} \\ i \sin \frac{\theta^c}{2} & \cos \frac{\theta^c}{2} \end{pmatrix} \begin{pmatrix} e^{\frac{i\psi^c}{2}} & 0 \\ 0 & e^{-\frac{i\psi^c}{2}} \end{pmatrix} \equiv \\ &\equiv \mathbf{g}(\varphi^c, 0, 0) \mathbf{g}(0, \theta^c, 0) \mathbf{g}(0, 0, \psi^c). \end{aligned}$$

Therefore,

$$\begin{aligned} \mathbf{g}(\varphi_1^c, \theta_1^c, \psi_1^c) \mathbf{g}(\varphi_2^c, \theta_2^c, \psi_2^c) &= \\ &\mathbf{g}(\varphi_1^c, 0, 0) \mathbf{g}(0, \theta_1^c, 0) \mathbf{g}(0, 0, \psi_1^c) \mathbf{g}(\varphi_2^c, 0, 0) \mathbf{g}(0, \theta_2^c, 0) \mathbf{g}(0, 0, \psi_2^c). \end{aligned} \quad (33)$$

It is obvious that

$$\mathbf{g}(0, 0, \psi_1^c) \mathbf{g}(\varphi_2^c, 0, 0) = \mathbf{g}(\varphi_2^c + \psi_1^c, 0, 0).$$

Besides, if we multiply the matrix $\mathbf{g}(\varphi^c, \theta^c, \psi^c)$ at the left by the matrix $\mathbf{g}(\varphi_1^c, 0, 0)$ the Euler angle φ^c increases by φ_1^c , and other Euler angles remain unaltered. Analogously, if we multiply at the right the matrix $\mathbf{g}(\varphi^c, \theta^c, \psi^c)$ by $\mathbf{g}(0, 0, \psi_2^c)$ the angle ψ^c increases by ψ_2^c . Hence it follows that in general case the angle φ_2^c should be replaced by $\varphi_2^c + \psi_1^c$, and the angles φ^c and ψ^c should be replaced by $\varphi^c - \varphi_1^c$ and $\psi^c - \psi_2^c$, that is,

$$\cos \theta^c = \cos \theta_1^c \cos \theta_2^c - \sin \theta_1^c \sin \theta_2^c \cos(\varphi_2^c + \psi_1^c), \quad (34)$$

$$e^{i(\varphi^c - \varphi_1^c)} = \frac{\sin \theta_1^c \cos \theta_2^c + \cos \theta_1^c \sin \theta_2^c \cos(\varphi_2^c + \psi_1^c) + i \sin \theta_2^c \sin(\varphi_2^c + \psi_1^c)}{\sin \theta^c}, \quad (35)$$

$$e^{\frac{i(\varphi^c + \psi^c - \varphi_1^c - \psi_2^c)}{2}} = \frac{\cos \frac{\theta_1^c}{2} \cos \frac{\theta_2^c}{2} e^{\frac{i(\varphi_2^c + \psi_1^c)}{2}} - \sin \frac{\theta_1^c}{2} \sin \frac{\theta_2^c}{2} e^{-\frac{i(\varphi_2^c + \psi_1^c)}{2}}}{\cos \frac{\theta^c}{2}}. \quad (36)$$

Addition Theorem for hyperspherical functions Z_{mn}^l follows from the relation

$$T_l(\mathbf{g}_1 \mathbf{g}_2) = T_l(\mathbf{g}_1) T_l(\mathbf{g}_2).$$

Hence it follows that

$$t_{mn}^l(\mathbf{g}_1 \mathbf{g}_2) = \sum_{k=-l}^l t_{mk}^l(\mathbf{g}_1) t_{kn}^l(\mathbf{g}_2).$$

Let us apply this equality to the matrices $\mathbf{g}_1$ and $\mathbf{g}_2$ with the Euler angles $0, 0, \theta_1, \tau_1, 0, 0$ and $\varphi_2, \epsilon_2, \theta_2, \tau_2, 0, 0$, correspondingly. Using the formula (24) we obtain

$$\begin{aligned} t_{mk}^l(\mathbf{g}_1) &= Z_{mk}^l(\cos \theta_1, \cosh \tau_1), \\ t_{kn}^l(\mathbf{g}_2) &= e^{-k(\epsilon_2 + i\varphi_2)} Z_{kn}^l(\cos \theta_2, \cosh \tau_2) \end{aligned}$$

and

where $\epsilon, \varphi, \theta, \tau, \varepsilon, \psi$ are the Euler angles of the matrix $\mathbf{g}_1 \mathbf{g}_2$. In accordance with (30)–(32) these angles are expressed via the factor angles $0, 0, \theta_1, \tau_1, 0, 0$ and $\varphi_2, \epsilon_2, \theta_2, \tau_2, 0, 0$. Thus, in this case the functions Z_{mn}^l satisfy the following addition theorem:

$$e^{-m(\epsilon+i\varphi)-n(\varepsilon+i\psi)} Z_{mn}^l(\cos \theta, \cosh \tau) = \sum_{k=-l}^l e^{-k(\epsilon_2+i\varphi_2)} Z_{mk}^l(\cos \theta_1, \cosh \tau_1) Z_{kn}^l(\cos \theta_2, \cosh \tau_2).$$

In general case when the Euler angles are related by the formulae (34)–(36) we obtain

$$e^{-m[\epsilon+\epsilon_1+i(\varphi_1-\varphi)]-n[\varepsilon+\varepsilon_2-i(\psi_2-\psi)]} Z_{mn}^l(\cos \theta, \cosh \tau) = \sum_{k=-l}^l e^{-k[\epsilon_2+\varepsilon_1+i(\varphi_2+\psi_1)]} Z_{mk}^l(\cos \theta_1, \cosh \tau_1) Z_{kn}^l(\cos \theta_2, \cosh \tau_2).$$

4 Infinitesimal operators of complex($SU(2)$) $\sim SL(2, \mathbb{C})$

Let $\omega^c(t)$ be the one-parameter subgroup of $SL(2, \mathbb{C})$. The operators of the right regular representation of $SL(2, \mathbb{C})$, corresponded to the elements of this subgroup, transfer complex functions $f(\mathbf{g})$ into $R(\omega^c(t))f(\mathbf{g}) = f(\mathbf{g}\omega^c(t))$. By this reason the infinitesimal operator of the right regular representation $R(\mathbf{g})$, associated with one-parameter subgroup $\omega^c(t)$, transfers the function $f(\mathbf{g})$ into $\frac{df(\mathbf{g}\omega^c(t))}{dt}$ at $t = 0$.

Let us denote Euler angles of the element $\mathbf{g}\omega^c(t)$ via $\varphi^c(t), \theta^c(t), \psi^c(t)$. Then there is an equality

$$\left. \frac{df(\mathbf{g}\omega^c(t))}{dt} \right|_{t=0} = \frac{\partial f}{\partial \varphi^c} (\varphi^c(0))' + \frac{\partial f}{\partial \theta^c} (\theta^c(0))' + \frac{\partial f}{\partial \psi^c} (\psi^c(0))'.$$

The infinitesimal operator A_ω^c , corresponded to the subgroup $\omega^c(t)$, has a form

$$A_\omega^c = A_\omega - iB_\omega = (\varphi^c(0))' \frac{\partial}{\partial \varphi^c} + (\theta^c(0))' \frac{\partial}{\partial \theta^c} + (\psi^c(0))' \frac{\partial}{\partial \psi^c}$$

Let us calculate infinitesimal operators A_1^c, A_2^c, A_3^c corresponding the complex subgroups $\Omega_1^c, \Omega_2^c, \Omega_3^c$. The subgroup Ω_3^c consists of the matrices

$$\omega_3(t^c) = \begin{pmatrix} e^{\frac{it^c}{2}} & 0 \\ 0 & e^{-\frac{it^c}{2}} \end{pmatrix}$$

Let $\mathbf{g} = \mathbf{g}(\varphi^c, \theta^c, \psi^c)$ be a matrix with complex Euler angles $\varphi^c = \varphi - i\epsilon$, $\theta^c = \theta - i\tau$, $\psi^c = \psi - i\varepsilon$. Therefore, Euler angles of the matrix $\mathbf{g}\omega_3(t^c)$ equal to $\varphi^c, \theta^c, \psi^c + t - it$. Hence it follows that

$$\varphi'(0) = 0, \quad \epsilon'(0) = 0, \quad \theta'(0) = 0, \quad \tau'(0) = 0, \quad \psi'(0) = 1, \quad \varepsilon'(0) = -i$$

So, the operator A_3^c , corresponded to the subgroup Ω_3^c , has a form

$$A_3^c = \frac{\partial}{\partial \psi^c} - i \frac{\partial}{\partial \varepsilon}$$

Whence

$$A_3 = \frac{\partial}{\partial \psi}, \quad (37)$$

$$B_3 = \frac{\partial}{\partial \varepsilon}. \quad (38)$$

Let us calculate the infinitesimal operator A_1^c corresponded the complex subgroup Ω_1^c . The subgroup Ω_1^c consists of the following matrices

$$\omega_1(t^c) = \begin{pmatrix} \cos \frac{t^c}{2} & i \sin \frac{t^c}{2} \\ i \sin \frac{t^c}{2} & \cos \frac{t^c}{2} \end{pmatrix}.$$

The Euler angles of these matrices equal to 0, $t^c = t - it$, 0. Let us represent the matrix $\mathbf{g}\omega_1(t^c)$ by the product $\mathbf{g}_1\mathbf{g}_2$, the Euler angles of which are described by the formulae (34)–(36). Then the Euler angles of the matrix $\omega_1(t^c)$ equal to $\varphi_2^c = 0$, $\theta_2^c = t - it$, $\psi_2^c = 0$, and the Euler angles of the matrix $\mathbf{g}$ equal to $\varphi_1^c = \varphi^c$, $\theta_1^c = \theta^c$, $\psi_1^c = \psi^c$. Thus, from the general formulae (34)–(36) we obtain that Euler angles $\varphi^c(t)$, $\theta^c(t)$, $\psi^c(t)$ of the matrix $\mathbf{g}\omega_1(t^c)$ are defined by the following relations:

$$\cos \theta^c(t) = \cos \theta^c \cos t^c - \sin \theta^c \sin t^c \cos \psi^c, \quad (39)$$

$$e^{i\varphi^c(t)} = e^{i\varphi^c} \frac{\sin \theta^c \cos t^c + \cos \theta^c \sin t^c \cos \psi^c + i \sin t^c \sin \psi^c}{\sin \theta^c(t)}, \quad (40)$$

$$e^{\frac{i[\varphi^c(t) + \psi^c(t)]}{2}} = e^{\frac{i\varphi^c}{2}} \frac{\cos \frac{\theta^c}{2} \cos \frac{t^c}{2} e^{\frac{i\psi^c}{2}} - \sin \frac{\theta^c}{2} \sin \frac{t^c}{2} e^{-\frac{i\psi^c}{2}}}{\cos \frac{\theta^c(t)}{2}}. \quad (41)$$

For calculation of derivatives $\varphi'(t)$, $\epsilon'(t)$, $\theta'(t)$, $\tau'(t)$, $\psi'(t)$, $\varepsilon'(t)$ at $t = 0$ we differentiate on t the both parts of the each equality from (39)–(41) and take $t = 0$. At this point we have $\varphi(0) = \varphi$, $\epsilon(0) = \epsilon$, $\theta(0) = \theta$, $\tau(0) = \tau$, $\psi(0) = \psi$, $\varepsilon(0) = \varepsilon$.

So, let us differentiate the both parts of (39). In the result we obtain

$$\theta'(t) - i\tau'(t) = (1 - i) \cos \psi^c.$$

Taking $t = 0$ we find that

$$\theta'(0) = \cos \psi^c, \quad \tau'(0) = \cos \psi^c.$$

Differentiating now the both parts of (40) we obtain

$$\varphi'(0) - i\epsilon'(0) = \frac{(1 - i) \sin \psi^c}{\sin \theta^c}.$$

Therefore,

$$\varphi'(0) = \frac{\sin \psi^c}{\sin \theta^c}, \quad \epsilon'(0) = \frac{\sin \psi^c}{\sin \theta^c}$$

Further, differentiating the both parts of (41) we find that

$$\psi'(0) - i\varepsilon'(0) = -(1+i) \cot \theta^c \sin \psi^c$$

and

$$\psi'(0) = -\cot \theta^c \sin \psi^c, \quad \varepsilon'(0) = -\cot \theta^c \sin \psi^c.$$

In such a way, we obtain the following infinitesimal operators:

$$A_1 = \cos \psi^c \frac{\partial}{\partial \theta} + \frac{\sin \psi^c}{\sin \theta^c} \frac{\partial}{\partial \varphi} - \cot \theta^c \sin \psi^c \frac{\partial}{\partial \psi}, \quad (42)$$

$$B_1 = \cos \psi^c \frac{\partial}{\partial \tau} + \frac{\sin \psi^c}{\sin \theta^c} \frac{\partial}{\partial \epsilon} - \cot \theta^c \sin \psi^c \frac{\partial}{\partial \epsilon}. \quad (43)$$

Let us calculate now an infinitesimal operator A_2^c corresponded to the complex subgroup Ω_2^c . The subgroup Ω_2^c consists of the following matrices

$$\omega_2(t^c) = \begin{pmatrix} \cos \frac{t^c}{2} & -\sin \frac{t^c}{2} \\ \sin \frac{t^c}{2} & \cos \frac{t^c}{2} \end{pmatrix},$$

where the Euler angles equal correspondingly to 0, $t^c = t - it$, 0. It is obvious that the matrix $\mathbf{g}\omega_2(t^c)$ can be represented by the product

$$\mathbf{g}_1 \mathbf{g}_2 = \begin{pmatrix} \cos \frac{\theta_1^c}{2} & i \sin \frac{\theta_1^c}{2} \\ i \sin \frac{\theta_1^c}{2} & \cos \frac{\theta_1^c}{2} \end{pmatrix} \begin{pmatrix} \cos \frac{\theta_2^c}{2} e^{\frac{i\varphi_2^c}{2}} & -\sin \frac{\theta_2^c}{2} e^{\frac{i\varphi_2^c}{2}} \\ \sin \frac{\theta_2^c}{2} e^{-\frac{i\varphi_2^c}{2}} & \cos \frac{\theta_2^c}{2} e^{-\frac{i\varphi_2^c}{2}} \end{pmatrix}.$$

Multiplying the matrices in the right part of this equality we obtain that Euler angles of the product $\mathbf{g}_1 \mathbf{g}_2$ are related by formulae

$$\cos \theta^c = \cos \theta_1^c \cos \theta_2^c + \sin \theta_1^c \sin \theta_2^c \sin \varphi_2^c, \quad (44)$$

$$e^{i\varphi^c} = \frac{\sin \theta_1^c \cos \theta_2^c - \cos \theta_1^c \sin \theta_2^c \sin \varphi_2^c + i \sin \theta_2^c \cos \varphi_2^c}{\sin \theta^c}, \quad (45)$$

$$e^{\frac{i(\varphi^c + \psi^c)}{2}} = \frac{\cos \frac{\theta_1^c}{2} \cos \frac{\theta_2^c}{2} e^{\frac{i\varphi_2^c}{2}} + i \sin \frac{\theta_1^c}{2} \sin \frac{\theta_2^c}{2} e^{-\frac{i\varphi_2^c}{2}}}{\cos \frac{\theta^c}{2}}. \quad (46)$$

Or, repeating the calculations as in the case of (34)–(36) we obtain in general case

$$\cos \theta^c = \cos \theta_1^c \cos \theta_2^c + \sin \theta_1^c \sin \theta_2^c \sin(\varphi_2^c + \psi_1^c), \quad (47)$$

$$e^{i(\varphi^c - \varphi_1^c)} = \frac{\sin \theta_1^c \cos \theta_2^c - \cos \theta_1^c \sin \theta_2^c \sin(\varphi_2^c + \psi_1^c) + i \sin \theta_2^c \cos(\varphi_2^c + \psi_1^c)}{\sin \theta^c}, \quad (48)$$

$$e^{\frac{i(\varphi^c + \psi^c - \varphi_1^c - \psi_2^c)}{2}} = \frac{\cos \frac{\theta_1^c}{2} \cos \frac{\theta_2^c}{2} e^{\frac{i(\varphi_2^c + \psi_1^c)}{2}} + \sin \frac{\theta_1^c}{2} \sin \frac{\theta_2^c}{2} e^{-\frac{i(\varphi_2^c + \psi_1^c)}{2}}}{\cos \frac{\theta^c}{2}}. \quad (49)$$

Therefore, Euler angles of the matrix $\omega_2(t^c)$ equal to $\varphi_2^c = 0$, $\theta_2^c = t - it$, $\psi_2^c = 0$, and Euler angles of the matrix $\mathbf{g}$ equal to $\varphi_1^c = \varphi^c$, $\theta_1^c = \theta^c$, $\psi_1^c = \psi^c$. Then from the formulae (47)–(49) we obtain that Euler angles $\varphi^c(t)$, $\theta^c(t)$, $\psi^c(t)$ of the matrix $\mathbf{g}\omega_2(t^c)$ are defined by relations

$$\cos \theta^c(t) = \cos \theta^c \cos t^c + \sin \theta^c \sin t^c \sin \psi^c, \quad (50)$$

$$e^{i\varphi^c(t)} = e^{i\varphi^c} \frac{\sin \theta^c \cos t^c - \cos \theta^c \sin t^c \sin \psi^c + i \sin t^c \cos \psi^c}{\sin \theta^c(t)}, \quad (51)$$

$$e^{\frac{i[\varphi^c(t) + \psi^c(t)]}{2}} = e^{\frac{i\varphi^c}{2}} \frac{\cos \frac{\theta^c}{2} \cos \frac{t^c}{2} e^{\frac{i\psi^c}{2}} + i \sin \frac{\theta^c}{2} \sin \frac{t^c}{2} e^{-\frac{i\psi^c}{2}}}{\cos \frac{\theta^c(t)}{2}}. \quad (52)$$

Differentiating on t the both parts of the each equalities (50)–(52) and taking $t = 0$ we obtain

$$\begin{aligned} \theta'(0) &= \tau'(0) = -\sin \psi^c, \\ \varphi'(0) &= \epsilon'(0) = \frac{\cos \psi^c}{\sin \theta^c}, \\ \psi'(0) &= \varepsilon'(0) = -\cot \theta^c \cos \psi^c. \end{aligned} \quad (53)$$

Therefore, for the subgroup Ω_2^c we have the following infinitesimal operators

$$A_2 = -\sin \psi^c \frac{\partial}{\partial \theta} + \frac{\cos \psi^c}{\sin \theta^c} \frac{\partial}{\partial \varphi} - \cot \theta^c \cos \psi^c \frac{\partial}{\partial \psi}, \quad (54)$$

$$B_2 = -\sin \psi^c \frac{\partial}{\partial \tau} + \frac{\cos \psi^c}{\sin \theta^c} \frac{\partial}{\partial \epsilon} - \cot \theta^c \cos \psi^c \frac{\partial}{\partial \varepsilon}. \quad (55)$$

It is easy to verify that operators A_i , B_i , defined by the formulae (37), (38), (42), (43) and (54), (55), are satisfy the commutation relations (1).

5 Recurrence relations between hyperspherical functions

Between generalized hyperspherical functions $\mathfrak{M}_{mn}^l$ (and also the hyperspherical functions Z_{mn}^l) there exists a wide variety of recurrence relations. Part of them relates the hyperspherical functions of one and the same order (with identical l), other part relates the functions of different orders.

In virtue of the Van der Waerden representation (5) the recurrence formulae for the hyperspherical functions of one and the same order follow from the equalities

$$X_- \mathfrak{M}_{mn}^l = \alpha'_n \mathfrak{M}_{m,n-1}^l, \quad X_+ \mathfrak{M}_{mn}^l = \alpha'_{n+1} \mathfrak{M}_{m,n+1}^l, \quad (56)$$

$$Y_- \mathfrak{M}_{mn}^l = \alpha_n \mathfrak{M}_{m,n-1}^l, \quad Y_+ \mathfrak{M}_{mn}^l = \alpha_{n+1} \mathfrak{M}_{m,n+1}^l, \quad (57)$$

where

$$\alpha'_n = \frac{\sqrt{(n+1)(n-m+1)}}{\sqrt{(n+1)(n-m+1)}}, \quad \alpha_n = \frac{\sqrt{(n+1)(n-m+1)}}{\sqrt{(n+1)(n-m+1)}}$$

From (2) and (4) it follows that

$$\begin{aligned} X_+ &= \frac{1}{2} (A_1 + iA_2 + iB_1 - B_2), \\ X_- &= \frac{1}{2} (A_1 - iA_2 + iB_1 + B_2), \\ Y_+ &= \frac{1}{2} (A_1 + iA_2 - iB_1 + B_2), \\ Y_- &= \frac{1}{2} (A_1 - iA_2 - iB_1 - B_2), \end{aligned}$$

Using the formulae (42), (43) and (54), (55) we obtain

$$X_+ = \frac{e^{-i\psi^c}}{2} \left[\frac{\partial}{\partial\theta} + \frac{i}{\sin\theta^c} \frac{\partial}{\partial\varphi} - i \cot\theta^c \frac{\partial}{\partial\psi} + i \frac{\partial}{\partial\tau} - \frac{1}{\sin\theta^c} \frac{\partial}{\partial\epsilon} + \cot\theta^c \frac{\partial}{\partial\epsilon} \right], \quad (58)$$

$$X_- = \frac{e^{i\psi^c}}{2} \left[\frac{\partial}{\partial\theta} - \frac{i}{\sin\theta^c} \frac{\partial}{\partial\varphi} + i \cot\theta^c \frac{\partial}{\partial\psi} + i \frac{\partial}{\partial\tau} + \frac{1}{\sin\theta^c} \frac{\partial}{\partial\epsilon} - \cot\theta^c \frac{\partial}{\partial\epsilon} \right], \quad (59)$$

$$Y_+ = \frac{e^{-i\psi^c}}{2} \left[\frac{\partial}{\partial\theta} + \frac{i}{\sin\theta^c} \frac{\partial}{\partial\varphi} - i \cot\theta^c \frac{\partial}{\partial\psi} - i \frac{\partial}{\partial\tau} + \frac{1}{\sin\theta^c} \frac{\partial}{\partial\epsilon} - \cot\theta^c \frac{\partial}{\partial\epsilon} \right], \quad (60)$$

$$Y_- = \frac{e^{i\psi^c}}{2} \left[\frac{\partial}{\partial\theta} - \frac{i}{\sin\theta^c} \frac{\partial}{\partial\varphi} + i \cot\theta^c \frac{\partial}{\partial\psi} - i \frac{\partial}{\partial\tau} - \frac{1}{\sin\theta^c} \frac{\partial}{\partial\epsilon} + \cot\theta^c \frac{\partial}{\partial\epsilon} \right]. \quad (61)$$

Further, substituting the function $\mathfrak{M}_{mn}^l = e^{-m(\epsilon-i\varphi)} Z_{mn}^l(\theta, \tau) e^{-n(\epsilon-i\psi)}$ into the relations (56) and taking into account the operators (58) and (59) we find that

$$\frac{\partial \dot{Z}_{mn}^l}{\partial\theta} + i \frac{\partial \dot{Z}_{mn}^l}{\partial\tau} - \frac{2(m-n\cos\theta^c)}{\sin\theta^c} \dot{Z}_{mn}^l = 2\alpha'_n \dot{Z}_{m,n-1}^l, \quad (62)$$

$$\frac{\partial \dot{Z}_{mn}^l}{\partial\theta} + i \frac{\partial \dot{Z}_{mn}^l}{\partial\tau} + \frac{2(m-n\cos\theta^c)}{\sin\theta^c} \dot{Z}_{mn}^l = 2\alpha'_{n+1} \dot{Z}_{m,n+1}^l. \quad (63)$$

Since the functions $\dot{Z}_{mn}^l(\theta, \tau)$ are symmetric, that is $\dot{Z}_{mn}^l(\theta, \tau) = \dot{Z}_{nm}^l(\theta, \tau)$, then substituting $\dot{Z}_{nm}^l(\theta, \tau)$ in lieu of $\dot{Z}_{mn}^l$ into the formulae (62)–(63) and replacing m by n , and n by m , we obtain

$$\frac{\partial \dot{Z}_{mn}^l}{\partial\theta} + i \frac{\partial \dot{Z}_{mn}^l}{\partial\tau} - \frac{2(n-m\cos\theta^c)}{\sin\theta^c} \dot{Z}_{mn}^l = 2\alpha'_m \dot{Z}_{m-1,n}^l, \quad (64)$$

$$\frac{\partial \dot{Z}_{mn}^l}{\partial\theta} + i \frac{\partial \dot{Z}_{mn}^l}{\partial\tau} + \frac{2(n-m\cos\theta^c)}{\sin\theta^c} \dot{Z}_{mn}^l = 2\alpha'_{m+1} \dot{Z}_{m+1,n}^l. \quad (65)$$

From (62)–(63) and (64)–(65) it follows that

$$\alpha'_{n+1} \dot{Z}_{m,n+1}^l - \alpha'_n \dot{Z}_{m,n-1}^l = \frac{2(m-n\cos\theta^c)}{\sin\theta^c} \dot{Z}_{mn}^l, \quad (66)$$

$$\alpha'_{m+1} \dot{Z}_{m+1,n}^l - \alpha'_m \dot{Z}_{m-1,n}^l = \frac{2(n-m\cos\theta^c)}{\sin\theta^c} \dot{Z}_{mn}^l. \quad (67)$$

Analogously, for the relations (57) we have

$$\frac{\partial Z_{mn}^l}{\partial\theta} - i \frac{\partial Z_{mn}^l}{\partial\tau} + \frac{2(m-n\cos\theta^c)}{\sin\theta^c} Z_{mn}^l = 2\alpha_n Z_{m,n-1}^l, \quad (68)$$

Further, using the symmetry of the functions Z_{mn}^l we obtain

$$\frac{\partial Z_{mn}^l}{\partial \theta} - i \frac{\partial Z_{mn}^l}{\partial \tau} + \frac{2(n - m \cos \theta^c)}{\sin \theta^c} Z_{mn}^l = 2\alpha_m Z_{m-1,n}^l, \quad (70)$$

$$\frac{\partial Z_{mn}^l}{\partial \theta} - i \frac{\partial Z_{mn}^l}{\partial \tau} - \frac{2(n - m \cos \theta^c)}{\sin \theta^c} Z_{mn}^l = 2\alpha_{m+1} Z_{m+1,n}^l. \quad (71)$$

Therefore,

$$\alpha_n Z_{m,n-1}^l - \alpha_{n+1} Z_{m,n+1}^l = \frac{2(m - n \cos \theta^c)}{\sin \theta^c} Z_{mn}^l, \quad (72)$$

$$\alpha_m Z_{m-1,n}^l - \alpha_{m+1} Z_{m+1,n}^l = \frac{2(n - m \cos \theta^c)}{\sin \theta^c} Z_{mn}^l. \quad (73)$$

Let us consider now recurrence relations between hyperspherical functions with different order. These recurrence formulae are related with the tensor products of irreducible representations of the Lorentz group. Indeed, in accordance with Van der Waerden representation (5) an arbitrary finite-dimensional representation of the group $\mathfrak{G}_+$ has a form $\tau_{l_0} \otimes \tau_{0l'} \sim \tau_{ll'}$, where τ_{l_0} and $\tau_{0l'}$ are representations of the group $SU(2)$. Then a product of the two representations $\tau_{l_1 l'_1}$ and $\tau_{l_2 l'_2}$ of the Lorentz group is defined by an expression

$$\tau_{l_1 l'_1} \otimes \tau_{l_2 l'_2} = \sum_{|l_1 - l_2| \leq k \leq l_1 + l_2; |l'_1 - l'_2| \leq k' \leq l'_1 + l'_2} \tau_{kk'}.$$

The vectors $e_{mm'}^{ll'}$ of the canonical basis have the form

$$e_{mm'}^{ll'} = \sum_{j+k=m, j'+k'=m'} C(l_1, l_2, l; j, k, m) C(l'_1, l'_2, l'; j', k', m') e_{jj'} \otimes e_{kk'}, \quad (74)$$

where

$$C(l_1, l_2, l; j, k, m) C(l'_1, l'_2, l'; j', k', m') = B_{l_1, l_2, l; l'_1, l'_2, l'}^{j, k, m; j', k', m'}$$

are the Clebsch–Gordan coefficients of the group $SL(2, \mathbb{C})$. Expressing the Clebsch–Gordan coefficients $C(l_1, l_2, l; j, k, j+k)$ of the group $SU(2)$ via a generalized hypergeometric function ${}_3F_2$ (see, for example [12, 23]) we see that CG-coefficients of $SL(2, \mathbb{C})$ have the form

$$\begin{aligned} B_{l_1, l_2, l; l'_1, l'_2, l'}^{j, k, m; j', k', m'} &= (-1)^{l_1 + l'_1 - j - j'} \times \\ &\quad \frac{\Gamma(l_1 + l_2 - m + 1) \Gamma(l'_1 + l'_2 - m' + 1)}{\Gamma(l_2 - l_1 + m + 1) \Gamma(l'_2 - l'_1 + m' + 1)} \times \\ &\quad \sqrt{\frac{\Gamma(l - m + 1) \Gamma(l + l_2 - l_1 + 1) \Gamma(l_1 - j + 1) \Gamma(l_2 + k + 1) \Gamma(l + m + 1) (2l + 1)}{\Gamma(l_1 - l_2 + l + 1) \Gamma(l_1 + l_2 - l + 1) \Gamma(l_1 + l_2 + l + 1) \Gamma(l_1 - j + 1) \Gamma(l_2 - k + 1)}} \times \\ &\quad \sqrt{\frac{\Gamma(l' - m' + 1) \Gamma(l' + l'_2 - l'_1 + 1) \Gamma(l'_1 - j' + 1) \Gamma(l'_2 + k' + 1) \Gamma(l' + m' + 1) (2l' + 1)}{\Gamma(l'_1 - l'_2 + l' + 1) \Gamma(l'_1 + l'_2 - l' + 1) \Gamma(l'_1 + l'_2 + l' + 1) \Gamma(l'_1 - j' + 1) \Gamma(l'_2 - k' + 1)}} \times \\ &\quad {}_3F_2((l + m + 1, -l + m, -l_1 + j; -l_1 - l_2 + m, l_2 - l_1 + m + 1; 1) \times \\ &\quad \frac{\Gamma((l'_1 + l'_2 - l' + 1) \Gamma(l'_1 + l'_2 - l' + 1) \Gamma(l'_1 + l'_2 + l' + 1) \Gamma(l'_1 - j' + 1) \Gamma(l'_2 - k' + 1)}{\Gamma(l'_1 - l'_2 + l' + 1) \Gamma(l'_1 + l'_2 - l' + 1) \Gamma(l'_1 + l'_2 + l' + 1) \Gamma(l'_1 - j' + 1) \Gamma(l'_2 - k' + 1)} \times \end{aligned} \quad (75)$$

where $m = j + k$, $m' = j' + k'$. In virtue of the orthogonality of the Clebsch–Gordan coefficients from (74) it follows that

$$\begin{aligned} e_{jj'} \otimes e_{kk'} &= \sum_{j+k=m, j'+k'=m'} \overline{C(l_1, l_2, l; j, k, m) C(l'_1, l'_2, l'; j', k', m')} e_{mm'}^{ll'} \\ &= \sum_{j+k=m, j'+k'=m'} \bar{B}_{l_1, l_2, l; l'_1, l'_2, l'}^{j, k, m; j', k', m'} e_{mm'}^{ll'}. \end{aligned} \quad (76)$$

Let us assume that $l = l'$ (this case is a most important for physics, since the Dirac field $(1/2, 0) \oplus (0, 1/2)$, the Weyl field $(1/2, 0) \cup (0, 1/2)$ and the Maxwell field $(1, 0) \cup (0, 1)$ are defined in terms of the functions (9) at $l = l'$). Therefore, at these restrictions we have $e_{jj'} \sim e_j$, $e_{kk'} \sim f_k$ and $e_{mm'}^{ll'} \sim g_m^l$. Further, assume that $l_1 = 1$ and $l_2 = l$, then the system (76) can be rewritten as follows

$$\begin{aligned} e_{-1} \otimes f_{m+1} &= \bar{b}_{11}^m g_m^{l+1} + \bar{b}_{12}^m g_m^l + \bar{b}_{13}^m g_m^{l-1}, \\ e_0 \otimes f_m &= \bar{b}_{21}^m g_m^{l+1} + \bar{b}_{22}^m g_m^l + \bar{b}_{23}^m g_m^{l-1}, \\ e_1 \otimes f_{m-1} &= \bar{b}_{31}^m g_m^{l+1} + \bar{b}_{32}^m g_m^l + \bar{b}_{33}^m g_m^{l-1}, \end{aligned} \quad (77)$$

where

$$\begin{aligned} \bar{b}_{ik}^m &= \begin{pmatrix} \bar{B}_{1, l_2, l_2-1; 1, l_2, l_2-1}^{1, k, m; 1, k, m} & \bar{B}_{1, l_2, l_2-1; 1, l_2, l_2-1}^{0, k, m; 0, k, m} & \bar{B}_{1, l_2, l_2-1; 1, l_2, l_2-1}^{-1, k, m; -1, k, m} \\ \bar{B}_{1, l_2, l_2; 1, l_2, l_2}^{1, k, m; 1, k, m} & \bar{B}_{1, l_2, l_2; 1, l_2, l_2}^{0, k, m; 0, k, m} & \bar{B}_{1, l_2, l_2; 1, l_2, l_2}^{-1, k, m; -1, k, m} \\ \bar{B}_{1, l_2, l_2+1; 1, l_2, l_2+1}^{1, k, m; 1, k, m} & \bar{B}_{1, l_2, l_2+1; 1, l_2, l_2+1}^{0, k, m; 0, k, m} & \bar{B}_{1, l_2, l_2+1; 1, l_2, l_2+1}^{-1, k, m; -1, k, m} \end{pmatrix} = \\ &= \begin{pmatrix} \frac{(l-m)(l-m+1)}{(2l+1)(2l+2)} & \frac{(l+m+1)(l-m)}{2l(l+1)} & \frac{(l+m)(l+m+1)}{2l(2l+1)} \\ \frac{(l+m+1)(l-m+1)}{(2l+1)(l+1)} & \frac{m^2}{l(l+1)} & \frac{(l+m)(l-m)}{l(2l+1)} \\ \frac{(l+m)(l+m+1)}{(2l+1)(2l+2)} & \frac{(l+m)(l-m+1)}{2l(l+1)} & \frac{(l-m)(l-m+1)}{2l(2l+1)} \end{pmatrix}. \end{aligned}$$

Let $T_{\mathfrak{g}}^l$ be a matrix of the irreducible representation of the weight l in the canonical basis. Let us apply the transformation $T_{\mathfrak{g}}$ to the left and right parts of the each equalities (77). In the left part we have

$$T_{\mathfrak{g}} e_k \otimes f_{m-k} = T_{\mathfrak{g}}^1 e_k \otimes T_{\mathfrak{g}}^l f_{m-k} = \bar{b}_{k+2,1}^m T_{\mathfrak{g}}^{l+1} g_m^{l+1} + \bar{b}_{k+2,2}^m T_{\mathfrak{g}}^l g_m^l + \bar{b}_{k+2,3}^m T_{\mathfrak{g}}^{l-1} g_m^{l-1},$$

where $k = -1, 0, 1$. Denoting the elements of $T_{\mathfrak{g}}^l$ via $\mathfrak{M}_{mn}^l$ (generalized hyperspherical functions) we find

$$\begin{aligned} (\mathfrak{M}_{-1,k}^1 e_{-1} + \mathfrak{M}_{0,k}^1 e_0 + \mathfrak{M}_{1,k}^1 e_1) \sum \mathfrak{M}_{j,m-k}^l f_j = \\ \sum (\bar{b}_{k+2,1}^m \mathfrak{M}_{jm}^{l+1} g_j^{l+1} + \bar{b}_{k+2,2}^m \mathfrak{M}_{jm}^l g_j^l + \bar{b}_{k+2,3}^m \mathfrak{M}_{jm}^{l-1} g_j^{l-1}). \end{aligned}$$

Replacing in the right part the vectors g_j^{l+1} , g_j^l , g_j^{l-1} via $e_{-1} \otimes f_{j+1}$, $e_0 \otimes f_j$, $e_1 \otimes f_{j-1}$ and comparing the coefficients at $e_{-1} \otimes f_{j+1}$, $e_0 \otimes f_j$, $e_1 \otimes f_{j-1}$ in the left and right parts we obtain three relations depending on k . Giving in these relations three possible values $-1, 0, 1$ to the number k and substituting the functions $\mathfrak{M}_{mn}^1(\varphi, \epsilon, \theta, \tau, \psi, \epsilon)$ (the matrix (29)) we find

the following nine recurrence relations:

$$\begin{aligned}
& \bar{b}_{11}^m \mathfrak{M}_{jm}^{l+1} \bar{b}_{11}^j + \bar{b}_{12}^m \mathfrak{M}_{jm}^l \bar{b}_{12}^j + \bar{b}_{13}^m \mathfrak{M}_{jm}^{l-1} \bar{b}_{13}^j = \\
& \quad \left(\cos^2 \frac{\theta}{2} \cosh^2 \frac{\tau}{2} + \frac{i \sin \theta \sinh \tau}{2} - \sin^2 \frac{\theta}{2} \sinh^2 \frac{\tau}{2} \right) e^{\epsilon+i\varphi+\epsilon+i\psi} \mathfrak{M}_{j+1,m+1}^l, \\
& \bar{b}_{11}^m \mathfrak{M}_{jm}^{l+1} \bar{b}_{21}^j + \bar{b}_{12}^m \mathfrak{M}_{jm}^l \bar{b}_{22}^j + \bar{b}_{13}^m \mathfrak{M}_{jm}^{l-1} \bar{b}_{23}^j = \frac{1}{\sqrt{2}} (\cos \theta \sinh \tau + i \sin \theta \cosh \tau) e^{-\epsilon-i\psi} \mathfrak{M}_{j,m+1}^l, \\
& \bar{b}_{11}^m \mathfrak{M}_{jm}^{l+1} \bar{b}_{31}^j + \bar{b}_{12}^m \mathfrak{M}_{jm}^l \bar{b}_{32}^j + \bar{b}_{13}^m \mathfrak{M}_{jm}^{l-1} \bar{b}_{33}^j = \\
& \quad \left(\cos^2 \frac{\theta}{2} \sinh^2 \frac{\tau}{2} + \frac{i \sin \theta \sinh \tau}{2} - \sin^2 \frac{\theta}{2} \cosh^2 \frac{\tau}{2} \right) e^{\epsilon+i\varphi-\epsilon-i\psi} \mathfrak{M}_{j-1,m+1}^l, \\
& \bar{b}_{21}^m \mathfrak{M}_{jm}^{l+1} \bar{b}_{11}^j + \bar{b}_{22}^m \mathfrak{M}_{jm}^l \bar{b}_{12}^j + \bar{b}_{23}^m \mathfrak{M}_{jm}^{l-1} \bar{b}_{13}^j = \frac{1}{\sqrt{2}} (\cos \theta \sinh \tau + i \sin \theta \cosh \tau) e^{\epsilon+i\psi} \mathfrak{M}_{j+1,m}^l, \\
& \bar{b}_{21}^m \mathfrak{M}_{jm}^{l+1} \bar{b}_{21}^j + \bar{b}_{22}^m \mathfrak{M}_{jm}^l \bar{b}_{22}^j + \bar{b}_{23}^m \mathfrak{M}_{jm}^{l-1} \bar{b}_{23}^j = (\cos \theta \cosh \tau + i \sin \theta \sinh \tau) \mathfrak{M}_{j,m}^l, \\
& \bar{b}_{21}^m \mathfrak{M}_{jm}^{l+1} \bar{b}_{31}^j + \bar{b}_{22}^m \mathfrak{M}_{jm}^l \bar{b}_{32}^j + \bar{b}_{23}^m \mathfrak{M}_{jm}^{l-1} \bar{b}_{33}^j = \frac{1}{\sqrt{2}} (\cos \theta \sinh \tau + i \sin \theta \cosh \tau) e^{-\epsilon-i\psi} \mathfrak{M}_{j-1,m}^l, \\
& \bar{b}_{31}^m \mathfrak{M}_{jm}^{l+1} \bar{b}_{11}^j + \bar{b}_{32}^m \mathfrak{M}_{jm}^l \bar{b}_{12}^j + \bar{b}_{33}^m \mathfrak{M}_{jm}^{l-1} \bar{b}_{13}^j = \\
& \quad \left(\cos^2 \frac{\theta}{2} \sinh^2 \frac{\tau}{2} + \frac{i \sin \theta \sinh \tau}{2} - \sin^2 \frac{\theta}{2} \cosh^2 \frac{\tau}{2} \right) e^{-\epsilon-i\varphi+\epsilon+i\psi} \mathfrak{M}_{j+1,m-1}^l, \\
& \bar{b}_{31}^m \mathfrak{M}_{jm}^{l+1} \bar{b}_{21}^j + \bar{b}_{32}^m \mathfrak{M}_{jm}^l \bar{b}_{22}^j + \bar{b}_{33}^m \mathfrak{M}_{jm}^{l-1} \bar{b}_{23}^j = \frac{1}{\sqrt{2}} (\cos \theta \sinh \tau + i \sin \theta \cosh \tau) e^{-\epsilon-i\varphi} \mathfrak{M}_{j,m-1}^l, \\
& \bar{b}_{31}^m \mathfrak{M}_{jm}^{l+1} \bar{b}_{31}^j + \bar{b}_{32}^m \mathfrak{M}_{jm}^l \bar{b}_{32}^j + \bar{b}_{33}^m \mathfrak{M}_{jm}^{l-1} \bar{b}_{33}^j = \\
& \quad \left(\cos^2 \frac{\theta}{2} \cosh^2 \frac{\tau}{2} + \frac{i \sin \theta \sinh \tau}{2} - \sin^2 \frac{\theta}{2} \sinh^2 \frac{\tau}{2} \right) e^{-\epsilon-i\varphi-\epsilon-i\psi} \mathfrak{M}_{j-1,m-1}^l.
\end{aligned}$$

Let us find recurrence relations between the functions $\mathfrak{M}_{mn}^l$, where the weight l changed by $\frac{1}{2}$. As known, the product of an irreducible representation with $l = \frac{1}{2}$ by an arbitrary irreducible representation with the weight l is decomposed into a representation with $l + \frac{1}{2}$ (a basis in the corresponded space has a form $g_m^{l+\frac{1}{2}}$ ($-l - \frac{1}{2} \leq m \leq l + \frac{1}{2}$) and a representation with $l - \frac{1}{2}$ ($g_m^{l-\frac{1}{2}}$ ($-l + \frac{1}{2} \leq m \leq l - \frac{1}{2}$)). Thus, by analogy with (77) we obtain

$$\begin{aligned}
e_{\frac{1}{2}} \otimes f_{m-\frac{1}{2}} &= \bar{b}_{00}^m g_m^{l+\frac{1}{2}} + \bar{b}_{01}^m g_m^{l-\frac{1}{2}}, \\
e_{-\frac{1}{2}} \otimes f_{m+\frac{1}{2}} &= \bar{b}_{10}^m g_m^{l+\frac{1}{2}} + \bar{b}_{11}^m g_m^{l-\frac{1}{2}},
\end{aligned}$$

where

$$\bar{b}_{ik} = \begin{pmatrix} \bar{B}_{\frac{1}{2}, l_2, l_2 - \frac{1}{2}; \frac{1}{2}, k, m} & \bar{B}_{\frac{1}{2}, l_2, l_2 - \frac{1}{2}; \frac{1}{2}, k, m} \\ \bar{B}_{\frac{1}{2}, l_2, l_2 + \frac{1}{2}; \frac{1}{2}, k, m} & \bar{B}_{\frac{1}{2}, l_2, l_2 + \frac{1}{2}; \frac{1}{2}, k, m} \end{pmatrix} = \begin{pmatrix} \frac{l_2 - m + \frac{1}{2}}{2l_2 + 1} & \frac{l_2 + m + \frac{1}{2}}{2l_2 + 1} \\ \frac{l_2 + m + \frac{1}{2}}{2l_2 + 1} & \frac{l_2 - m + \frac{1}{2}}{2l_2 + 1} \end{pmatrix}.$$

Carrying out the analogous calculations as for the case $l = 1$ and using the matrix (28) we

come to the following recurrence relations

$$\begin{aligned}
\bar{b}_{00}^m \mathfrak{M}_{jm}^{l+\frac{1}{2}} \bar{b}_{00}^j + \bar{b}_{01}^m \mathfrak{M}_{jm}^{l-\frac{1}{2}} \bar{b}_{01}^j &= \left(\cos \frac{\theta}{2} \cosh \frac{\tau}{2} + i \sin \frac{\theta}{2} \sinh \frac{\tau}{2} \right) e^{\frac{\epsilon+i\varphi+\varepsilon+i\psi}{2}} \mathfrak{M}_{j+\frac{1}{2}, m+\frac{1}{2}}^l, \\
\bar{b}_{00}^m \mathfrak{M}_{jm}^{l+\frac{1}{2}} \bar{b}_{10}^j + \bar{b}_{01}^m \mathfrak{M}_{jm}^{l-\frac{1}{2}} \bar{b}_{11}^j &= \left(\cos \frac{\theta}{2} \sinh \frac{\tau}{2} + i \sin \frac{\theta}{2} \cosh \frac{\tau}{2} \right) e^{\frac{\epsilon+i\varphi-\varepsilon-i\psi}{2}} \mathfrak{M}_{j-\frac{1}{2}, m+\frac{1}{2}}^l, \\
\bar{b}_{10}^m \mathfrak{M}_{jm}^{l+\frac{1}{2}} \bar{b}_{00}^j + \bar{b}_{11}^m \mathfrak{M}_{jm}^{l-\frac{1}{2}} \bar{b}_{01}^j &= \left(\cos \frac{\theta}{2} \sinh \frac{\tau}{2} + i \sin \frac{\theta}{2} \cosh \frac{\tau}{2} \right) e^{\frac{-\epsilon-i\varphi+\varepsilon+i\psi}{2}} \mathfrak{M}_{j+\frac{1}{2}, m-\frac{1}{2}}^l, \\
\bar{b}_{10}^m \mathfrak{M}_{jm}^{l+\frac{1}{2}} \bar{b}_{10}^j + \bar{b}_{11}^m \mathfrak{M}_{jm}^{l-\frac{1}{2}} \bar{b}_{11}^j &= \left(\cos \frac{\theta}{2} \cosh \frac{\tau}{2} + i \sin \frac{\theta}{2} \sinh \frac{\tau}{2} \right) e^{\frac{-\epsilon-i\varphi-\varepsilon-i\psi}{2}} \mathfrak{M}_{j-\frac{1}{2}, m-\frac{1}{2}}^l.
\end{aligned}$$

References

- [1] A.I. Ahiezer, V.B. Berestetskii, Quantum Electrodynamics (Nauka, Moscow, 1969) [in Russian].
- [2] D.V. Ahluwalia, D.J. Ernst, $(j, 0) \oplus (0, j)$ covariant spinors and causal propagators based on Weinberg formalism, Int. J. Mod. Phys. **E2**, 397–422 (1993).
- [3] P. Appell, M.J. Kampé de Fériet, Fonctions hypégeométriques et hypersphériques. Polynomes d’Hermite (Gauthier–Villars, 1926).
- [4] A.O. Barut, R. Raczka, Theory of Group Representations and Applications (PWN, Warszawa, 1977).
- [5] V.V. Dvoeglazov, *Extra Dirac equations*, Nuovo Cimento, **B111**, 483–496 (1996).
- [6] A. Erdélyi, Higher Transcendental Functions, vol. I (Mc Grow-Hill Book Company, New York, 1953).
- [7] I.M. Gel’fand, Z.Ya. Shapiro, it Representations of the rotation group of three-dimensional space and their applications, Uspekhi Mat. Nauk **7**, 3–117 (1952) [in Russian].
- [8] I.M. Gel’fand, R.A. Minlos, Z.Ya. Shapiro, Representations of the Rotation and Lorentz Groups and their Applications (Pergamon Press, Oxford, 1963).
- [9] D.M. Gitman, A.L. Shelepin, *Fields on the Poincaré Group: Arbitrary Spin Description and Relativistic Wave Equations*, Int. J. Theor. Phys. **40**(3), 603–684 (2001).
- [10] G. Mackey, Induced Representations of Groups and Quantum Mechanics (Benjamin, New York, 1968).
- [11] M.A. Naimark, Linear Representations of the Lorentz Group (Pergamon, London, 1964).
- [12] M.E. Rose, Multipole Fields (John Wiley & Sons, New York, 1955).
- [13] Yu.B. Rumer, A.I. Fet, Group Theory and Quantized Fields (Nauka, Moscow, 1977) [in Russian].
- [14] L. Ryder, Quantum Field Theory (Cambridge University Press, Cambridge, 1985).
- [15] S.S. Schweber, An Introduction to Relativistic Quantum Field Theory (Harper & Row, New

- [16] V.V. Varlamov, *Fundamental Automorphisms of Clifford Algebras and an Extension of Dąbrowski Pin Groups*, Hadronic J. **22**, 497–535 (1999).
- [17] V.V. Varlamov, *Discrete Symmetries and Clifford Algebras*, Int. J. Theor. Phys. **40**(4), 769–805 (2001).
- [18] V.V. Varlamov, *Clifford Algebras and Lorentz Group*, preprint math-ph/0108022 (2001).
- [19] V.V. Varlamov, *About Algebraic Foundations of Majorana-Oppenheimer Quantum Electrodynamics and de Broglie-Jordan Neutrino Theory of Light*, preprint math-ph/0109024; to appear in special issue “Contemporary Electrodynamics” of the journal “Annales de la Fondation de Louis de Broglie”.
- [20] V.V. Varlamov, *Group Theoretical Description of Space Inversion, Time Reversal and Charge Conjugation*, preprint math-ph/0203059 (2002).
- [21] M.A. Vasiliev, *Higher-spin gauge theories in four, three and two dimensions*, Int. J. Mod. Phys. D**5**, 763–797 (1996).
- [22] N.Ya. Vilenkin, *Special Functions and the Theory of Group Representations* (AMS, Providence, 1968).
- [23] N.Ya. Vilenkin, A.U. Klimyk, *Representations of Lie Groups and Special Functions*, vols. 1–3. (Dordrecht: Kluwer Acad. Publ., 1991–1993).
- [24] B.L. van der Waerden, *Die Gruppentheoretische Methode in der Quantenmechanik* (Springer, Berlin, 1932).
- [25] S. Weinberg, *Feinman rules for any spin I & II & III*, Phys. Rev. **133B**, 1318–1332 & **134B**, 882–896 (1964) & **181B**, 1893–1899 (1969).
- [26] E.P. Wigner, *On unitary representations of the inhomogeneous Lorentz group*, Ann. Math. **40**, 149–204 (1939).

Department of Mathematics
 Siberia State University of Industry
 Kirova 42, Novokuznetsk 654007
 Russia